

Extensions of C^* -dynamical systems to systems with complete transfer operators

B.K. Kwaśniewski

Institute of Mathematics, University of Białystok,
ul. Akademicka 2, PL-15-267 Białystok, Poland
e-mail: bartoszk@math.uwb.edu.pl
<http://math.uwb.edu.pl/~zaf/kwasniewski>

Abstract

Starting from an arbitrary endomorphism δ of a unital C^* -algebra $\mathcal{A}$ we construct a bigger algebra $\mathcal{B}$ and extend δ onto $\mathcal{B}$ in such a way that $\delta : \mathcal{B} \rightarrow \mathcal{B}$ has a unital kernel and a hereditary range, i.e. there exists a unique non-degenerate transfer operator for $(\mathcal{B}, \delta)$. The pair $(\mathcal{B}, \delta)$ is universal with respect to a suitable notion of covariant representation.

Keywords: C^* -algebra, endomorphism, partial isometry, crossed product, covariant representation, transfer operator

2000 Mathematics Subject Classification: 46L55, 46L30, 16W20

Contents

1	Covariant representations	3
2	Transfer operators.	7
3	Extensions of C^*-dynamical systems up to systems with complete transfer operators	10
3.1	Adjoining the unit to the kernel of an endomorphism	11
3.2	Inductive limit construction	13
4	Commutative case and its relation to inverse limits	17
4.1	Compactification of a complement of a range of a mapping	18
4.2	Inverse limits and spectral description	19
4.3	Comparison of the spectral and the algebraic approach	22

Introduction

The crossed-product of a C^* -algebra $\mathcal{A}$ by an automorphism $\delta : \mathcal{A} \rightarrow \mathcal{A}$ is defined as a universal C^* -algebra generated by a copy of $\mathcal{A}$ and a unitary element U satisfying the relations

$$\delta(a) = UaU^*, \quad \delta^{-1}(a) = U^*aU, \quad a \in \mathcal{A}.$$

Algebras arising in this way (or their versions adapted to actions of groups of automorphisms) are very well understood and became one of the standard constructions in C^* -theory, see for instance [KR86], [Ped79]. On the other hand, the natural desire to adapt this kind of constructions to endomorphisms (or semigroups of endomorphisms) encounters serious obstacles from the very beginning. Roughly speaking, it is caused by the *irreversibility* of the system $(\mathcal{A}, \delta)$ (the lack of δ^{-1}) in case δ is an endomorphism, and the main question is what relations should the element U satisfy.

The difficulty of the matter manifests itself in a variety of approaches, see, for example, [CK80], [Pas80], [Sta93], [Mur96], [Exel94], [Exel03], [Kwa05a], [ABL05] which do however have a certain nontrivial intersection. They mostly agree, and simultaneously boast their greatest successes, in the case when dynamics is implemented by monomorphisms with a hereditary range. In view of the recent article [BL05], see also [ABL05], [Kwa07], this coincidence is completely understood. It is shown in [BL05] that in the case of monomorphism with hereditary range there exists a unique non-degenerate transfer operator δ_* for $(\mathcal{A}, \delta)$, called by authors of [BL05] a *complete transfer operator*, and the theory goes smooth with δ_* as it takes over the role classically played by δ^{-1} .

The goal of the present paper is an interesting fact that starting from an arbitrary endomorphism δ of an arbitrary unital C^* -algebra $\mathcal{A}$ there is a universal (from a point of view to be explained later) procedure of extending the C^* -dynamical system $(\mathcal{A}, \delta)$ to a bigger system $(\mathcal{B}, \delta)$ possessing a complete transfer operator. Philosophically, it is a procedure of passing from an irreversible dynamical system to a bigger reversible one. In commutative case it takes on a literal meaning as then the C^* -dynamical systems $(\mathcal{A}, \delta)$ and $(\mathcal{B}, \delta)$ correspond to topological dynamical systems (X, α) and $(\tilde{X}, \tilde{\alpha})$ respectively. Chronologically this topological approach was the first one that led to the description of $(\mathcal{B}, \delta)$, see [KL03], [Kwa05a]. The relationship between (X, α) and $(\tilde{X}, \tilde{\alpha})$ is of particular interest. For instance, the space $\tilde{X}$ may be viewed as a generalization of the topological inverse limit space, see [Kwa05a], [Kwa05b], and Section 4 of the present paper. Thus as a rule $\tilde{X}$ is topologically very complicated - contains indecomposable continua [Nad92], has a structure of hyperbolic attractors [Wil67], [Wil70], or of a space arising from substitution tilings [AP98].

The paper is organized as follows.

In the first section we introduce two notions of a covariant representation of a C^* -dynamical system $(\mathcal{A}, \delta)$. The first one exists in a literature in a similar or identical form, see, for example, [LO04], [Exel03], [Mur96], [Sta93]. The second one which we shall call a *strict covariant representation* is more restrictive, however, it seems to reconcile most of the previous versions of this kind of notion. After a closer look this former definition becomes completely natural and it is essential for our purposes. Second section serves as a collection of definitions and facts concerning transfer operators for C^* -dynamical systems. It is based on [Kwa07] and [BL05], and the only new result there is Proposition 2.6 which states that for C^* -dynamical systems possessing a complete transfer operator strict covariant representations defined in

this paper agrees with covariant representations defined in [BL05].

The third section is the core of the article. It contains the main result the first technical step of which is an extension of the endomorphism $\delta : \mathcal{A} \rightarrow \mathcal{A}$ to a one with unital kernel. Next for systems $(\mathcal{A}, \delta)$ where $\ker \delta$ is unital we define algebra $\mathcal{B}$ as a direct limit of a certain direct limit of C^* -algebras

$$\mathcal{B}_0 \xrightarrow{\delta_0} \mathcal{B}_1 \xrightarrow{\delta_1} \mathcal{B}_2 \xrightarrow{\delta_2} \dots$$

The form of this sequence will become relatively clear after analyzing the case of the system acting on a Hilbert space, cf. Proposition 3.1. It is worth to mention here that the first attempt to such analysis was made in [LO04]. Algebra $\mathcal{A}$ embeds into $\mathcal{B}$ and extension of δ onto $\mathcal{B}$ is induced on $\mathcal{B}$ by an inverse limit. Distinctive property of the system $(\mathcal{B}, \delta)$ is that it possess a complete transfer operator δ_* and in fact it might be thought as a universal extension of $(\mathcal{A}, \delta)$ with respect to this property and strict covariant representations.

The closing section is devoted to commutative (or one would say topological) systems. In particular, we discuss the relationship between inverse limit spaces and the spectrum of algebra $\mathcal{B}$, we present a slight generalization of a main result from [KL03] and set it against the general approach from Section 3.

1 Covariant representations

Throughout this paper we let $\mathcal{A}$ be a C^* -algebra with an identity 1 and $\delta : \mathcal{A} \rightarrow \mathcal{A}$ be an endomorphism of this algebra (all the morphisms appearing in the text are assumed to be $*$ -preserving). We shall refer to the pair $(\mathcal{A}, \delta)$ as a C^* -dynamical system.

In order to introduce and clarify the role of the relations we shall use in the present article, let us start with a non-trivial Theorem that gathers information on the subject developed by different authors. Formally it should be placed at the end of Section 3 because a part of it is based on Theorem 3.7 to be proved there. However, as it reveals the underlying associations the author decided to place it here.

By a *covariant triple* for a C^* -algebra $\mathcal{A}$ we will mean any triple (π, U, H) consisting of a non-degenerate faithful representation $\pi : \mathcal{A} \rightarrow \mathcal{B}(H)$ on a Hilbert space H and a partial isometry $U \in \mathcal{B}(H)$.

Theorem 1.1. *Let $\delta, \delta_* : \mathcal{A} \rightarrow \mathcal{A}$ be two mappings on a unital C^* -algebra $\mathcal{A}$. Consider the following conditions that a covariant triple (π, U, H) may satisfy*

$$(CR1) \quad U\pi(a) = \pi(\delta(a))U, \quad a \in \mathcal{A},$$

$$(CR2) \quad \pi(\delta(a)) = U\pi(a)U^*, \quad a \in \mathcal{A},$$

$$(CR3) \quad \pi(\delta_*(a)) = U^*\pi(a)U, \quad a \in \mathcal{A}.$$

Then

- a) $\delta : \mathcal{A} \rightarrow \mathcal{A}$ is an endomorphism of $\mathcal{A}$ if and only if there exists a covariant triple (π, U, H) satisfying (CR1) and (CR2),

- b) $\delta_* : \mathcal{A} \rightarrow \mathcal{A}$ is a non-degenerate transfer operator for an endomorphism δ (see Definition 2.1) if and only if there exists a covariant triple (π, U, H) satisfying (CR1) and (CR3),
- c) $\delta_* : \mathcal{A} \rightarrow \mathcal{A}$ is a complete transfer operator for an endomorphism δ (see Definition 2.3) if and only if there exists a covariant triple (π, U, H) satisfying (CR1), (CR2) and (CR3).

Item a) is a corollary to the main result of the present paper, see Theorem 3.7. Item b) follows from [Exel03, Theorem 3.4] and item c) is the main result of [BL05].

A definition of a covariant representation for $(\mathcal{A}, \delta)$ is naturally determined by Theorem 1.1.

Definition 1.2. Let $(\mathcal{A}, \delta)$ be a C^* -dynamical system. A covariant triple (π, U, H) for $\mathcal{A}$ satisfying (CR1) and (CR2) will be called a *covariant representation* of $(\mathcal{A}, \delta)$.

Another useful condition which in the presence of (CR2) may be used interchangeably with (CR1) is

$$(\mathbf{CR1}') \quad U^*U \in \pi(\mathcal{A})',$$

where $\pi(\mathcal{A})'$ is a commutant of $\pi(\mathcal{A})$ in $\mathcal{B}(H)$, see [LO04, Proposition 2.2].

Proposition 1.3. For any mapping $\delta : \mathcal{A} \rightarrow \mathcal{A}$ and any covariant triple (π, U, H) satisfying (CR2) conditions (CR1) and (CR1') are equivalent.

Apart from being relatively natural Definition 1.2 has a certain disadvantage. Namely, a projection U^*U might not reflect the structure of the kernel of endomorphism δ . For instance, one would like to have the following implications: if δ is a monomorphism (an automorphism) then U is an isometry (a unitary operator respectively), however condition (CR1) (equivalently (CR1')) does not ensure this.

Example 1.4. Let $\mathcal{A} \subset \mathcal{B}(H)$ and $\delta(\cdot) = U(\cdot)U^*$ where $U \in \mathcal{B}(H)$ is a partial isometry such that $U^*U \in Z(\mathcal{A})$. Then the algebra $\mathcal{A}$ is a direct sum

$$\mathcal{A} = (1 - U^*U)\mathcal{A} \oplus U^*U\mathcal{A}$$

where $(1 - U^*U)\mathcal{A} = \ker \delta$ is the kernel of δ . Let us define a representation $\pi : \mathcal{A} \rightarrow \mathcal{B}(\tilde{H})$ where $\tilde{H} = \bigoplus_{n=0}^{\infty} H$ and a partial isometry $\tilde{U} \in \mathcal{B}(\tilde{H})$ by the formulae

$$(\pi(a)h)_n = ah_n, \quad a \in \mathcal{A}, \quad (\tilde{U}h)_n = Uh_{n+1}, \quad n \in \mathbb{N}.$$

Then $(\pi, \tilde{U}, \tilde{H})$ is a covariant representation of $(\mathcal{A}, \delta)$. However, the operator $\tilde{U}$ is never an isometry because

$$(\tilde{U}^*\tilde{U}h)_n = \begin{cases} 0 & , \quad n = 0 \\ U^*Uh_n & , \quad n > 0 \end{cases},$$

and the algebra $(1 - \tilde{U}^*\tilde{U})\pi(\mathcal{A}) \cong \mathcal{A}$ is not related to $(1 - U^*U)\mathcal{A} = \ker \delta$ at all. In particular, we have

$$\tilde{U}^*\tilde{U} < \pi(U^*U).$$

It turns out that the obstacle from Example 1.4 illustrate the general rule.

Proposition 1.5. *Let $(\mathcal{A}, \delta)$ be a C^* -dynamical system and (π, U, H) a covariant triple satisfying (CR2). Let $P \in \mathcal{B}(H)$ be an orthogonal projection given by the limit in the strong operator topology*

$$P = 1 - s \lim_{\lambda \in \Lambda} \pi(\mu_\lambda), \quad (1)$$

where $\{\mu_\lambda\}_{\lambda \in \Lambda}$ is an approximate unit for $\ker \delta$. Then

$$U^*U \leq P \quad \text{and} \quad P \in \pi(\mathcal{A})'.$$

Moreover, if $U^*U \in \pi(\mathcal{A})$, then the kernel of $\delta : \mathcal{A} \rightarrow \mathcal{A}$ is unital and $U^*U = P$ is the complement of the unit in $\pi(\ker \delta)$.

Proof. Firstly let us pass to the algebra $\mathcal{A}_1 = C^*(\pi(\mathcal{A}), P)$ generated by $\pi(\mathcal{A})$ and the projection P . Since $\ker \delta$ is an ideal in $\mathcal{A}$, $P \in \pi(\mathcal{A})'$ and thus $\mathcal{A}_1 = P\pi(\mathcal{A}) \oplus (1 - P)\pi(\mathcal{A})$. Moreover, the mapping $\tilde{\delta}(\cdot) = U(\cdot)U^*$ is an endomorphism of $\mathcal{A}_1$ (extending the initial endomorphism $\pi\delta : \pi(\mathcal{A}) \rightarrow \pi(\mathcal{A})$). Indeed, one readily checks that for $a, b \in \mathcal{A}$ we have

$$\tilde{\delta}(P\pi(a) + (1 - P)\pi(b)) = U(P\pi(a) + (1 - P)\pi(b))U^* = \pi(\delta(a)).$$

Let $a \in \mathcal{A}_1$. We assert that the following statements are equivalent

- i) $Pa = 0$
- ii) $U^*UaU^*U = 0$
- iii) $U^*Ua = 0$
- iv) $aU^*U = 0$

Equivalence $i) \Leftrightarrow ii)$ follows from the fact that $P\pi(\mathcal{A}) \cong \pi(\delta(\mathcal{A}))$ and $\pi(\delta(\mathcal{A})) \cong U^*U\pi(\mathcal{A})U^*U$ where the isomorphisms are given by the correspondences $P\pi(a) \longleftrightarrow \pi(\delta(a))$ and $\pi(\delta(a)) \longleftrightarrow U^*U\pi(a)U^*U$, respectively. Implications $iii) \Rightarrow ii)$, $iv) \Rightarrow ii)$ are obvious. To show $ii) \Rightarrow iii)$, $ii) \Rightarrow iv)$ we note that $\{b \in \mathcal{A}_1 : U^*UbU^*U = 0\} = (1 - P)\pi(\mathcal{A})$ is an ideal in $\mathcal{A}_1$ and thus if $U^*UaU^*U = 0$, then

$$\|U^*Ua\|^2 = \|U^*Ua(U^*Ua)^*\| + \|U^*U(aa^*)U^*U\| = 0,$$

$$\|aU^*U\|^2 = \|(aU^*U)^*aU^*U\| + \|U^*U(a^*a)U^*U\| = 0.$$

Hence our assertion is true. In particular, $U^*U(1 - P) = 0$ and $U^*U \leq P$ holds. Moreover, if we assume that $U^*U \in \pi(\mathcal{A})$ then $1 - U^*U$ belongs to the kernel of $\delta : \mathcal{A}_1 \rightarrow \mathcal{A}_1$ and, in view of the assertion we proved above, it is a unit in this kernel. Hence $1 - U^*U = 1 - P$ and thus $U^*U = P \in \pi(\mathcal{A})$ and $\mathcal{A}_1 = \pi(\mathcal{A})$. $\blacksquare$

The extreme case in the above proposition, that is when $U^*U = P$, plays an essential role in the present paper. Hence we introduce one more condition:

(CR1'') $U^*U = 1 - s \lim_{\lambda \in \Lambda} \pi(\mu_\lambda)$

where $\{\mu_\lambda\}_{\lambda \in \Lambda}$ is an approximate unit in $\ker \delta$.

In view of Example 1.4 the conditions (CR1') and (CR1'') are not equivalent but Proposition 1.5 shows that, in the presence of (CR2), (CR1'') implies (CR1').

Definition 1.6. A covariant triple (π, U, H) for $\mathcal{A}$ satisfying (CR1'') and (CR2) will be called a *strict covariant representation* of a C^* -dynamical system $(\mathcal{A}, \delta)$.

Every strict covariant representation is a covariant representation and one of the corollaries to the main result of the present paper is that any C^* -dynamical system possess a strict covariant representation. We have the following immediate corollaries to Proposition 1.5.

Corollary 1.7. *Let (π, U, H) be a covariant representation of $(\mathcal{A}, \delta)$ and let $\ker \delta$ be unital. The following conditions are equivalent*

- i) (π, U, H) is a strict covariant representation,
- ii) $U^*U \in \pi(\mathcal{A})$,
- iii) $U^*U \in \pi(Z(\mathcal{A}))$,
- iv) $1 - U^*U$ is the unit in $\pi(\ker \delta)$.

Corollary 1.8. *For any strict covariant representation (π, U, H) of a C^* -dynamical system $(\mathcal{A}, \delta)$ we have*

- i) δ is a monomorphism $\iff U$ is an isometry,
- ii) δ is an automorphism $\implies U$ is unitary.

We finish this subsection showing that every covariantly represented system $(\mathcal{A}, \delta)$ may be extended to a system strictly covariantly represented. What is important is that during this process of extension the only thing that may undergo an enlargement is the kernel of δ . First we formulate a straightforward but also very useful proposition.

Proposition 1.9. *For every covariant representation (π, U, H) of $(\mathcal{A}, \delta)$ the mapping*

$$U^*U\pi(a) \longmapsto \delta(a)$$

establishes an isomorphism $U^*U\pi(\mathcal{A}) \cong \delta(\mathcal{A})$. In particular the set

$$\{a \in \mathcal{A} : (1 - U^*U)\pi(a) = \pi(a)\}$$

coincide with the kernel of δ .

Proof. The mentioned isomorphism is given by $\pi^{-1}(U(\cdot)U^*)$ and its inverse is implemented by $U^*(\pi(\cdot))U$. Thus the second part of statement holds because we have

$$\delta(a) = 0 \iff U^*U\pi(a) = 0 \iff (1 - U^*U)\pi(a) = \pi(a).$$

■

Proposition 1.10. *Let (π, U, H) be a covariant representation of $(\mathcal{A}, \delta)$ and let P be given by (1). Let us put $\mathcal{A}_1 = C^*(\pi(\mathcal{A}), P)$ and $\mathcal{A}_2 = C^*(\pi(\mathcal{A}), U^*U)$. Then*

$$U\mathcal{A}_iU^* \subset \mathcal{A}_i, \quad U^*U \in \mathcal{A}'_i, \quad i = 1, 2$$

and the C^* -dynamical systems $(\mathcal{A}_1, \tilde{\delta})$, $(\mathcal{A}_2, \tilde{\delta})$, where $\tilde{\delta}(\cdot) = U(\cdot)U^*$, are conjugate by a homomorphism $T = T_1 \oplus T_2$ which action is given by the diagram

$$\begin{array}{ccc} \mathcal{A}_2 & = & U^*U\pi(\mathcal{A}) \oplus (1 - U^*U)\pi(\mathcal{A}) \\ & \downarrow T_1 & \downarrow T_2 \\ \mathcal{A}_1 & = & P\pi(\mathcal{A}) \oplus (1 - P)\pi(\mathcal{A}) \end{array}$$

T_1 is an isomorphism and T_2 is an epimorphism. More precisely we have

$$T_1(U^*U\pi(a)) = P\pi(a), \quad T_2((1 - U^*U)\pi(a)) = (1 - P)\pi(a), \quad a \in \mathcal{A}.$$

Proof. In view of Proposition 1.5 we have $1 - U^*U \geq 1 - P$ and thereby the formula

$$T_2(\pi(a)(1 - U^*U)) = \pi(a)(1 - P), \quad a \in \mathcal{A},$$

defines an epimorphism $T_2 : \pi(\mathcal{A})(1 - U^*U) \rightarrow \pi(\mathcal{A})(1 - P)$.

If $a \in \mathcal{A}$ is such that $U^*Ua = 0$ then by Proposition 1.9, a is in the kernel of δ and hence $Pa = 0$. Thus the mapping T_1 is well defined and since $U^*U \leq P$ it is injective. Obviously T_1 is an isomorphism. The equality $\tilde{\delta} \circ T = T \circ \tilde{\delta}$ is checked readily. Namely, we have

$$\tilde{\delta}(T(U^*U\pi(a) + (1 - U^*U)\pi(b))) = T(\tilde{\delta}(U^*U\pi(a) + (1 - U^*U)\pi(b))) = \pi(\delta(a)).$$

■

Remark 1.11. The foregoing propositions give us a criterion for the existence of covariant representations. Namely, if we have a faithful nondegenerate representation $\pi : \mathcal{A} \rightarrow H$ and P is given by (1), then a necessary condition for existence of $U \in L(H)$ such that (π, U, H) is a covariant representation of $(\mathcal{A}, \delta)$, is that an ideal

$$I = \{a \in \mathcal{A} : (1 - P)\pi(a) = \pi(a)\}$$

coincides with the kernel of $\delta : \mathcal{A} \rightarrow \mathcal{A}$. Moreover, if this is the case then it is readily checked that an ideal

$$I^\perp = \{a \in \mathcal{A} : P\pi(a) = \pi(a)\}$$

is the biggest among all the ideals in $\mathcal{A}$ having a zero intersection with I .

2 Transfer operators.

In this section we list certain definitions and facts concerning transfer operators that are necessary for our future purposes. An inquiring reader is referred to [Kwa07], [BL05], [Exel03]. Everything henceforth up to Proposition 2.6 is borrowed from [Kwa07] and [BL05].

Definition 2.1. A linear map $\delta_*: \mathcal{A} \rightarrow \mathcal{A}$ is called a *transfer operator* for the pair $(\mathcal{A}, \delta)$ if it is continuous and positive and such that

$$\delta_*(\delta(a)b) = a\delta_*(b), \quad a, b \in \mathcal{A}. \quad (2)$$

If additionally one (and hence all) of the following equivalent conditions are satisfied

- i) the composition $E = \delta \circ \delta_*$ is a conditional expectation onto $\delta(\mathcal{A})$,
- ii) $\delta \circ \delta_* \circ \delta = \delta$,
- iii) $\delta(\delta_*(1)) = \delta(1)$,

then δ_* is called a *non-degenerate transfer operator*.

The following proposition presents conditions for existence and clarifies the structure of non-degenerate transfer operators.

Proposition 2.2. *There exists a non-degenerate transfer operator for $(\mathcal{A}, \delta)$ if and only if $\ker \delta$ is unital and there exists a conditional expectation $E: \delta(1)\mathcal{A}\delta(1) \rightarrow \delta(\mathcal{A})$.*

Moreover, then $1 - \delta_*(1)$ is the unit in $\ker \delta$ and we have

$$E(a) = \delta(\delta_*(a)), \quad \delta_*(a) = \delta^{-1}(E(\delta(1)a\delta(1))), \quad a \in \mathcal{A},$$

where δ^{-1} is the inverse to the isomorphism $\delta: \delta_*(1)\mathcal{A} \rightarrow \delta(\mathcal{A})$.

It follows that if $\delta(\mathcal{A}) = \delta(1)\mathcal{A}\delta(1)$, then there is a unique non-degenerate transfer operator for $(\mathcal{A}, \delta)$ (since there is a unique conditional expectation $E = id$). We recall [Exel03, Proposition 4.1] that the equality $\delta(\mathcal{A}) = \delta(1)\mathcal{A}\delta(1)$ holds if and only if $\delta(\mathcal{A})$ is a hereditary subalgebra of $\mathcal{A}$.

Definition 2.3. We shall say $(\mathcal{A}, \delta)$ is a *complete C^* -dynamical system* if $\delta: \mathcal{A} \rightarrow \mathcal{A}$ has a unital kernel and a hereditary range. Then, in view of Proposition 2.2, there is a unique non-degenerate transfer operator for $(\mathcal{A}, \delta)$ given by

$$\delta_*(a) = \delta^{-1}(\delta(1)a\delta(1)), \quad a \in \mathcal{A},$$

where δ^{-1} is the inverse to the isomorphism $\delta: (1 - q)\mathcal{A} \rightarrow \delta(\mathcal{A})$, q is the unit in $\ker \delta$. We shall say that δ_* is a *complete transfer operator* for $(\mathcal{A}, \delta)$.

The next result presents the criteria for the system to be complete.

Theorem 2.4. *Let $(\mathcal{A}, \delta)$ be a C^* -dynamical system. The following are equivalent:*

- i) the system $(\mathcal{A}, \delta)$ is complete,
- ii) there exists a transfer operator δ_* for $(\mathcal{A}, \delta)$ such that

$$\delta\delta_*(a) = \delta(1)a\delta(1), \quad a \in \mathcal{A}. \quad (3)$$

iii) there exists a non-degenerate transfer operator δ_* for $(\mathcal{A}, \delta)$ and $\delta(\mathcal{A})$ is a hereditary subalgebra of $\mathcal{A}$,

iv) there exists a central orthogonal projection $p \in \mathcal{A}$ such that the mapping $\delta : p\mathcal{A} \rightarrow \delta(\mathcal{A})$ is a $*$ -isomorphism, and $\delta(\mathcal{A}) = \delta(1)\mathcal{A}\delta(1)$,

Transfer operators in ii), iii) are equal (to the complete transfer operator for $(\mathcal{A}, \delta)$) and $p = \delta_*(1) = 1 - q$ where p is a projection from iv) and q is a unit for $\ker \delta$.

The article [BL05] provide us with a number of criteria implying the uniqueness of a non-degenerate transfer operator. In particular, it may be expressed in terms of covariant representations. More precisely, if δ_* is a non-degenerate transfer operator for $(\mathcal{A}, \delta)$ and (π, U, H) is a strict covariant representation then by Propositions we have

$$\pi(\delta_*(1)) = U^*U, \quad a \in \mathcal{A},$$

and one would wonder whether (CR3) is true. It follows from [BL05] that (CR3) holds only if δ_* is a complete transfer operator for $(\mathcal{A}, \delta)$.

Theorem 2.5 ([BL05], Theorem 3.1). *The system $(\mathcal{A}, \delta)$ is complete with a complete transfer operator δ_* if and only if there exist a covariant triple (π, U, H) satisfying conditions (CR1), (CR2) and (CR3).*

The next result shows that for complete systems the class of covariant triples from Theorem 2.5 coincide with the class of strict covariant representations.

Proposition 2.6. *Let $(\mathcal{A}, \delta)$ be a complete C^* -dynamical system with a complete transfer operator δ_* . Then a covariant representation (π, U, H) of $(\mathcal{A}, \delta)$ is strict if and only if it satisfies (CR3).*

Proof. If (π, U, H) satisfies (CR3) then $1 - U^*U = 1 - \pi(\delta_*(1))$ is the identity in $\pi(\ker \delta)$ and thus the covariant representation (π, U, H) is strict.

On the other hand, if we assume that (π, U, H) is strict, then $U(\cdot)U^* : U^*U\pi(\mathcal{A}) \rightarrow \pi(\delta(\mathcal{A}))$ is an isomorphism where the inverse one is given by $U^*(\cdot)U$ because

$$U^*\pi(\delta(a))U = U^*U\pi(a)U^*U = U^*U\pi(a), \quad a \in \mathcal{A}.$$

Furthermore, $U^*U = \pi(\delta_*(1))$ and $\delta(\mathcal{A}) = \delta(1)\mathcal{A}\delta(1)$. Thus, by the definition of a complete transfer operator we get

$$\begin{aligned} U^*\pi(a)U &= U^*(UU^*\pi(a)UU^*)U = U^*\pi(\delta(1)a\delta(1))U \\ &= U^*U\pi(\delta^{-1}(\delta(1)a\delta(1))) = \pi(\delta_*(1))\pi(\delta_*(a)) = \pi(\delta_*(a)), \quad a \in \mathcal{A}, \end{aligned}$$

what finishes the proof. $\blacksquare$

Corollary 2.7. *Any complete C^* -dynamical system possess a strict covariant representation.*

3 Extensions of C^* -dynamical systems up to systems with complete transfer operators

Before attempting the abstract construction it is reasonable and instructive to study the spatial case. We start with consideration of this sort what will result in establishing a basic structure of the extended algebra in Proposition 3.1.

Let us consider covariant representation of a C^* -dynamical system, that is let $U \in \mathcal{B}(H)$ and $\mathcal{A} \subset \mathcal{B}(H)$, $1 \in \mathcal{A}$, be such that

$$U\mathcal{A}U^* \subset \mathcal{A}, \quad U^*U \in \mathcal{A}'.$$

We have two operators acting on the C^* -algebra $\mathcal{B}(H)$, namely $\delta(\cdot) = U(\cdot)U^*$ and $\delta_*(\cdot) = U^*(\cdot)U$. Operator $\delta : \mathcal{A} \rightarrow \mathcal{A}$ is a $*$ -endomorphism however δ_* in general does not preserve $\mathcal{A}$. The situation changes when we pass to the C^* -algebra

$$\mathcal{B} = C^*\left(\bigcup_{n=0}^{\infty} U^{n*}\mathcal{A}U^n\right)$$

generated by $\bigcup_{n=0}^{\infty} U^{n*}\mathcal{A}U^n$. Then, see [LO04, Proposition 3.9], the following relations hold

$$UBU^* \subset \mathcal{B}, \quad U^*BU \subset \mathcal{B}, \quad U^*U \in Z(\mathcal{B}).$$

Thus the C^* -dynamical system $(\mathcal{B}, \delta)$ is an extension of the system $(\mathcal{A}, \delta)$ and furthermore operator $\delta_* : \mathcal{B} \rightarrow \mathcal{B}$ is a complete transfer operator for $(\mathcal{B}, \delta)$.

Let us define algebras "approximating" algebra $\mathcal{B}$ as C^* -algebras generated by finite sums:

$$\mathcal{B}_n = C^*\left(\bigcup_{i=0}^n U^{*i}\mathcal{A}U^i\right), \quad n \in \mathbb{N}.$$

The family $\{\mathcal{B}_n\}_{n \in \mathbb{N}}$ fixes the structure of a direct limit on $\mathcal{B}$. Indeed, we have

$$\mathcal{A} = \mathcal{B}_0 \subset \mathcal{B}_1 \subset \dots \subset \mathcal{B}_n \subset \dots, \quad \text{and} \quad \mathcal{B} = \overline{\bigcup_{n \in \mathbb{N}} \mathcal{B}_n}.$$

The coming proposition describes algebras $\mathcal{B}_n$ as direct sums of subalgebras of the algebra $C^*(\mathcal{A}, U^*U)$. In particular, if $U^*U \in \mathcal{A}$, then we are able to define algebras $\mathcal{B}_n$ in terms of the system $(\mathcal{A}, \delta)$.

Proposition 3.1. *Every element $a \in \mathcal{B}_n$ can be presented in the form*

$$a = a_0 + U^*a_1U + \dots + U^{*n}a_nU^n$$

where

$$a_i \in (1 - U^*U)\delta^i(1)\mathcal{A}\delta^i(1), \quad i = 0, \dots, n-1, \quad a_n \in \delta^n(1)\mathcal{A}\delta^n(1), \quad (4)$$

and this form is unique (if n is fixed). In particular, $a \mapsto a_0 \oplus a_1 \oplus \dots \oplus a_n$ establishes an isomorphism.

$$\mathcal{B}_n \cong (1 - U^*U)\mathcal{A} \oplus (1 - U^*U)\delta^1(1)\mathcal{A}\delta^1(1) \oplus \dots \oplus \delta^n(1)\mathcal{A}\delta^n(1)$$

Proof. Let $a \in \mathcal{B}_n$. Then $a = \sum_{k=0}^n \delta_*^k(b_k)$ where $b_i \in \mathcal{A}$ and $\delta_*(\cdot) = U^*(\cdot)U$, see [LO04, Proposition ??]. Without loss of generality we may assume that $b_i \in \delta^i(1)\mathcal{A}\delta^i(1)$, because $\delta_*^i(b_i) = \delta_*^i(\delta^i(1)b_i\delta^i(1))$. We shall construct elements a_i satisfying (4) modifying inductively the elements b_i .

For a_0 we take $b_0(1 - \delta_*(1))$ and "the remaining part" $b_0\delta_*(1)$ of b_0 we include in b_1 , that is we put $c_1 = b_1 + \delta(b_0)$. Then $a = a_0 + \delta_*(c_1) + \dots + \delta_*^n(b_n)$, because $b_0\delta_*(1) = \delta_*(\delta(b_0))$, and $a_0 \in (1 - U^*U)\mathcal{A}$.

Continuing in this manner we get $k < n$ coefficients $a_0, \dots, a_{k-1}$ satisfying (4) and such that $a = a_0 + \dots + \delta_*^{k-1}(a_{k-1}) + \delta_*^k(c_k) + \dots + \delta_*^n(b_n)$. We put $a_k = c_k(1 - \delta_*(1)) \in \mathcal{A}$ and $c_{k+1} = b_{k+1} + \delta(c_k)$. Then $a_k \in (1 - U^*U)\delta^k(1)\mathcal{A}\delta^k(1)$ whereas the following computation shows that $a = a_0 + \dots + \delta_*^k(a_k) + \delta_*^{k+1}(c_{k+1}) + \dots + \delta_*^n(b_n)$:

$$\begin{aligned} \delta_*^k(c_k) &= \delta_*^k(c_k)\delta_*^k(1) = \delta_*^k(c_k)(\delta_*^k(1) - \delta_*^{k+1}(1)) + \delta_*^{k+1}(1)\delta_*^k(c_k) \\ &= \delta_*^k(c_k(1 - \delta_*(1))) + \delta_*^{k+1}(\delta(c_k)) = \delta_*^k(a_k) + \delta_*^{k+1}(\delta(c_k)). \end{aligned}$$

This ends the first part of the proof.

To realize the consequences of conditions (4) we notice that they imply the following

$$\delta_*^i(a_i) \in (\delta_*^i(1) - \delta_*^{i+1}(1))\mathcal{A}, \quad i = 0, \dots, n-1. \quad (5)$$

Since the sequence $\{\delta_*^k(1)\}_{k \in \mathbb{N}} \subset Z(\mathcal{B})$ is a decreasing family of orthogonal projections we see that projections $1 - \delta_*(1)$, $\delta_*(1) - \delta_*^2(1)$, $\dots$, $\delta_*^n(1) - \delta_*^{n+1}(1)$, $\delta_*^n(1)$ are pair-wise orthogonal and central in $\mathcal{B}_n$. Hence the algebra $\mathcal{B}_n$ is a direct sum of ideals corresponding to these projections and i -th component of such decomposition is isomorphic to $(1 - U^*U)\delta^i(1)\mathcal{A}\delta^i(1)$, for $i = 0, \dots, n-1$ and $\delta^n(1)\mathcal{A}\delta^n(1)$ for $i = n$. $\blacksquare$

3.1 Adjoining the unit to the kernel of an endomorphism

Using Proposition 1.10 one may deduce that (up to an isomorphism) the algebra $C^*(\pi(\mathcal{A}), U^*U)$ does not depend on the choice of a strict covariant representation (π, U, H) of the system $(\mathcal{A}, \delta)$. In this subsection we present an "internal" (not involving covariant representations and hence independent from their existence) description of this algebra. We shall denote it by $\mathcal{A}^+$ and we shall consider endomorphism extension $\delta : \mathcal{A}^+ \rightarrow \mathcal{A}^+$ of δ up to $\mathcal{A}^+$ ($\mathcal{A} \subset \mathcal{A}^+$).

We shall construct algebra $\mathcal{A}^+$ by a general procedure of adjoining a unit to a given ideal in $\mathcal{A}$ (by an ideal we always mean two-sided closed ideal). Indeed, for any ideal I in $\mathcal{A}$ there exists the largest among the ideals J in $\mathcal{A}$ satisfying

$$I \cap J = \{0\}$$

(such an ideal could be defined directly as $(p\mathcal{A}^{**}) \cap \mathcal{A}$, see notation below). We denote it by $I^\perp$ and define $\mathcal{A}^+$ to be the following direct sum of quotient algebras

$$\mathcal{A}^+ = (\mathcal{A}/I) \oplus (\mathcal{A}/I^\perp).$$

The C^* -algebra $\mathcal{A}$ embeds naturally into C^* -algebra $\mathcal{A}^+$ via

$$\mathcal{A} \ni a \longmapsto (a + I) \oplus (a + I^\perp) \in \mathcal{A}^+.$$

Since $I \cap I^\perp = \{0\}$ this mapping is injective and we shall treat $\mathcal{A}$ as the corresponding subalgebra of $\mathcal{A}^+$. If we let I to be the kernel of endomorphism $\delta : \mathcal{A} \rightarrow \mathcal{A}$ then we can extend δ up to $\mathcal{A}^+$ by the formula

$$\mathcal{A}^+ \ni ((a + I) \oplus (b + I^\perp)) \xrightarrow{\delta} (\delta(a) + I) \oplus (\delta(b) + I^\perp) \in \mathcal{A}^+.$$

Since an element $\delta(a)$ does not depend on the choice of a representative of the equivalence class $a + I$ endomorphism $\delta : \mathcal{A}^+ \rightarrow \mathcal{A}^+$ is well defined (and obviously extends $\delta : \mathcal{A} \rightarrow \mathcal{A}$).

The fundamental property of the system $(\mathcal{A}^+, \delta)$ is that the kernel of the endomorphism $\delta : \mathcal{A}^+ \rightarrow \mathcal{A}^+$ is unital where its unit have the form $(0 + I) \oplus (1 + I^\perp)$. Evidently, if the kernel of $\delta : \mathcal{A} \rightarrow \mathcal{A}$ is unital the algebras $\mathcal{A}^+$ and $\mathcal{A}$ coincide.

We now present "an exterior" version of the preceding construction. It reveals universal character of $(\mathcal{A}^+, \delta)$ and will give us a desired correspondence between covariant representations of $(\mathcal{A}, \delta)$ and $(\mathcal{A}^+, \delta)$.

For that purpose, let us consider $\mathcal{A}$ as a subalgebra of $\mathcal{A}^{**}$ - the second dual algebra (enveloping W^* -algebra), and let $\{\mu_\lambda\}_{\lambda \in \Lambda}$ be an approximate unit for I . Then the σ -weak limits

$$q = \sigma \lim_{\lambda \in \Lambda} \mu_\lambda, \quad p = 1 - \sigma \lim_{\lambda \in \Lambda} \mu_\lambda$$

give rise to (pair-wise complementary) central projections in $\mathcal{A}^{**}$. Furthermore, one checks, cf. [Sak71], [KR86], that

$$I = (q\mathcal{A}^{**}) \cap \mathcal{A}, \quad I^\perp = (p\mathcal{A}^{**}) \cap \mathcal{A},$$

and consequently

$$\mathcal{A}/I \cong (\mathcal{A} + q\mathcal{A}^{**})/(q\mathcal{A}^{**}) \cong p\mathcal{A}, \quad \mathcal{A}/I^\perp \cong (\mathcal{A} + p\mathcal{A}^{**})/(p\mathcal{A}^{**}) \cong q\mathcal{A}.$$

Thus the algebra $\mathcal{A}^+$ may be viewed as a C^* -subalgebra of $\mathcal{A}^{**}$ generated by the algebra $\mathcal{A}$ and the projection p :

$$\mathcal{A}^+ = p\mathcal{A} + (1 - p)\mathcal{A} \subset \mathcal{A}^{**}.$$

Under this identification $\delta : \mathcal{A}^+ \rightarrow \mathcal{A}^+$ assumes the following form

$$\delta(pa + (1 - p)b) = \delta(a), \quad a, b \in \mathcal{A},$$

and the universality of $\mathcal{A}^{**}$ gives us the following statement.

Theorem 3.2. *There is a one-to-one correspondence between the strict covariant representations (π, U, H) and (π^+, U, H) of $(\mathcal{A}, \delta)$ and $(\mathcal{A}^+, \delta)$ respectively, which is established by the relation*

$$\pi^+((a + I) \oplus (b + I^\perp)) = \pi^+(pa + (1 - p)b) = U^*U\pi(a) + (1 - U^*U)\pi(b). \quad (6)$$

In particular, for every strict covariant representation (π, U, H) of $(\mathcal{A}, \delta)$ the algebra $\mathcal{A}^+$ is isomorphic to $C^*(U^*U, \pi(\mathcal{A}))$.

Proof. Representation π extends uniquely to a normal representation π'' of $\mathcal{A}^{**}$. We put $\pi^+ = \pi''|_{\mathcal{A}^+}$. Then by the strictness of (π, U, H) we have $\pi^+(p) = U^*U$ and thus π^+ is given by (6). In view of Remark 1.11 we have

$$I = \{a \in \mathcal{A} : U^*U\pi(a) = 0\}, \quad \text{and} \quad I^\perp = \{a \in \mathcal{A} : (1 - U^*U)\pi(a) = 0\},$$

what implies that $\pi^+|_{p\mathcal{A}}$ and $\pi^+|_{(1-p)\mathcal{A}}$ are injective. Thus π^+ is faithful and now it is easily seen that (π^+, U, H) is a strict covariant representation of $(\mathcal{A}^+, \delta)$. ■

3.2 Inductive limit construction

Now we are ready to pass to the main part of the construction. Heuristically it relies upon Proposition 3.1. For convenience, until Definition 3.5 we shall assume that the kernel of $\delta : \mathcal{A} \rightarrow \mathcal{A}$ is unital (in other case one may pass to the system $(\mathcal{A}^+, \delta)$ from the previous subsection). The unit in $\ker \delta$ will be denoted by q and its complementary projection by $p = 1 - q$. Thus q and p are central projections in $\mathcal{A}$.

We put

$$\mathcal{A}_n = \delta^n(1)\mathcal{A}\delta^n(1), \quad n \in \mathbb{N},$$

and define algebras $\mathcal{B}_n$ as direct sums of the form

$$\mathcal{B}_n = q\mathcal{A}_0 \oplus q\mathcal{A}_1 \oplus \dots \oplus q\mathcal{A}_{n-1} \oplus \mathcal{A}_n, \quad n \in \mathbb{N}. \quad (7)$$

In particular, $\mathcal{B}_0 = \mathcal{A}_0 = \mathcal{A}$.

Let us consider, for $n \in \mathbb{N}$, a homomorphism $\delta_n : \mathcal{B}_n \rightarrow \mathcal{B}_{n+1}$ schematically presented by the diagram

$$\begin{array}{ccccccc} \mathcal{B}_n & = & q\mathcal{A}_0 & \oplus & \dots & \oplus & q\mathcal{A}_{n-1} \oplus \mathcal{A}_n \\ \downarrow \delta_n & & \downarrow id & & & & \downarrow id \\ \mathcal{B}_{n+1} & = & q\mathcal{A}_0 & \oplus & \dots & \oplus & q\mathcal{A}_{n-1} \oplus q\mathcal{A}_n \oplus \mathcal{A}_{n+1} \end{array}$$

$\swarrow \delta$

and formally given by the formula

$$\delta_n(a_0 \oplus \dots \oplus a_{n-1} \oplus a_n) = a_0 \oplus \dots \oplus a_{n-1} \oplus qa_n \oplus \delta(a_n),$$

where $a_k \in q\mathcal{A}_k$, $k = 0, \dots, n-1$, and $a_n \in \mathcal{A}_n$. Homomorphism δ_n is well defined since $a_n = qa_n + pa_n$ and $\delta : p\mathcal{A} \rightarrow \delta(\mathcal{A})$ is an isomorphism δ_n is injective.

We define an algebra $\mathcal{B} = \varinjlim \{\mathcal{B}_n, \delta_n\}$ as a direct limit of the direct sequence

$$\mathcal{B}_0 \xrightarrow{\delta_0} \mathcal{B}_1 \xrightarrow{\delta_1} \mathcal{B}_2 \xrightarrow{\delta_2} \dots \quad (8)$$

and denote by $\phi_n : \mathcal{B}_n \rightarrow \mathcal{B}$, $n \in \mathbb{N}$, the natural embeddings, see for instance [Mur90]. Then

$$\phi_0(\mathcal{A}) = \phi_0(\mathcal{B}_0) \subset \phi_1(\mathcal{B}_1) \subset \dots \subset \phi_n(\mathcal{B}_n) \subset \dots \quad \text{and} \quad \mathcal{B} = \overline{\bigcup_{n \in \mathbb{N}} \phi_n(\mathcal{B}_n)}.$$

We shall identify algebra $\mathcal{A}$ with the subalgebra $\phi_0(\mathcal{A}) \subset \mathcal{B}$ and under this identification we extend δ onto the algebra $\mathcal{B}$. To this end, let us consider two sequences (an inverse one and a direct one)

$$\mathcal{B}_0 \xleftarrow{s_1} \mathcal{B}_1 \xleftarrow{s_2} \mathcal{B}_2 \xleftarrow{s_3} \dots, \quad (9)$$

$$\mathcal{B}_0 \xrightarrow{s_{*,0}} \mathcal{B}_1 \xrightarrow{s_{*,1}} \mathcal{B}_2 \xrightarrow{s_{*,2}} \dots, \quad (10)$$

where s_n is a "left-shift" and $s_{*,n}$ is a "right-shift":

$$s_n(a_0 \oplus a_1 \oplus \dots \oplus a_n) = a_1 \oplus a_2 \oplus \dots \oplus a_n$$

$$s_{*,n}(a_0 \oplus \dots \oplus a_{n-1} \oplus a_n) = 0 \oplus (\delta(1)a_0 \delta(1)) \oplus \dots \oplus (\delta^{n+1}(1)a_n \delta^{n+1}(1)),$$

$a_k \in q\mathcal{A}_k$, $k = 0, \dots, n-1$, $a_n \in \mathcal{A}_n$. Since $\delta^n(1)$, $n \in \mathbb{N}$, form a decreasing sequence of orthogonal projections, mappings s_n and $s_{*,n}$ are well defined. Moreover the operators s_n are homomorphisms, whereas operators $s_{*,n}$ may fail to be multiplicative.

Theorem 3.3. *Sequence (9) induces an endomorphism $\delta : \mathcal{B} \rightarrow \mathcal{B}$ extending the endomorphism $\delta : \mathcal{A} \rightarrow \mathcal{A}$, whereas sequence (10) induces an operator $\delta_* : \mathcal{B} \rightarrow \mathcal{B}$ which is a complete transfer operator for the extended C^* -dynamical system $(\mathcal{B}, \delta)$.*

The word "induces" means here that δ and δ_* are given on the dense $*$ -subalgebra $\bigcup_{n \in \mathbb{N}} \phi_n(\mathcal{A}_n)$ of $\mathcal{B}$ by the formulae

$$\delta(a) = \phi_n(s_{n+1}(\phi_{n+1}^{-1}(a))), \quad \delta_*(a) = \phi_{n+1}(s_{*,n}(\phi_n^{-1}(a))), \quad (11)$$

where $a \in \phi_n(\mathcal{B}_n)$, $n \in \mathbb{N}$.

Proof. Direct computations shows that the following diagrams

$$\begin{array}{ccccccc} \mathcal{B}_0 & \xrightarrow{\delta_0} & \mathcal{B}_1 & \xrightarrow{\delta_1} & \mathcal{B}_2 & \xrightarrow{\delta_2} & \dots & \xrightarrow{\delta_{n-1}} & \mathcal{B}_n & \xrightarrow{\delta_n} & \dots \\ & \nwarrow s_1 & & \nwarrow s_2 & & \nwarrow s_3 & & \nwarrow s_n & & \nwarrow s_{n+1} & \\ \mathcal{B}_0 & \xrightarrow{\delta_0} & \mathcal{B}_1 & \xrightarrow{\delta_1} & \mathcal{B}_2 & \xrightarrow{\delta_2} & \dots & \xrightarrow{\delta_{n-1}} & \mathcal{B}_n & \xrightarrow{\delta_n} & \dots \end{array}$$

$$\begin{array}{ccccccc} \mathcal{B}_0 & \xrightarrow{\delta_0} & \mathcal{B}_1 & \xrightarrow{\delta_1} & \mathcal{B}_2 & \xrightarrow{\delta_2} & \dots & \xrightarrow{\delta_{n-1}} & \mathcal{B}_n & \xrightarrow{\delta_n} & \dots \\ s_{*,0} \nearrow & & s_{*,1} \nearrow & & s_{*,2} \nearrow & & s_{*,n-1} \nearrow & & s_{*,n} \nearrow & & \\ \mathcal{B}_0 & \xrightarrow{\delta_0} & \mathcal{B}_1 & \xrightarrow{\delta_1} & \mathcal{B}_2 & \xrightarrow{\delta_2} & \dots & \xrightarrow{\delta_{n-1}} & \mathcal{B}_n & \xrightarrow{\delta_n} & \dots \end{array}$$

commute. Hence (9) and (10) induce certain linear mappings on $\mathcal{B}$ (i.e. formulae (11) make sense). The former mapping is a homomorphism (since s_n is a homomorphism for all $n \in \mathbb{N}$) and the latter one is positive (because $s_{*,n}$ posses that property for all $n \in \mathbb{N}$).

We assert that the mapping induced by (9) agrees with δ on $\mathcal{A}$ (which we identify with $\phi_0(\mathcal{A})$). Indeed, an element $a \in \phi_0(\mathcal{A})$ of the inductive limit $\mathcal{B}$ is represented by the following sequence

$$(a, qa \oplus \delta(a), qa \oplus q\delta(a) \oplus \delta^2(a), \dots)$$

which is mapped both by δ and the mapping induced by (9) onto the following sequence

$$(\delta(a), q\delta(a) \oplus \delta^2(a), q\delta(a) \oplus q\delta^2(a) \oplus \delta^3(a), \dots) \in \phi_0(\mathcal{A}).$$

Thereby our assertion is true and we are justified to denote by δ the mapping induced by (9).

By Theorem 2.4 to prove that δ_* is the complete transfer operator for $(\mathcal{B}, \delta)$ it suffices to show (2) and (3). For that purpose take arbitrary elements $\tilde{a}, \tilde{b} \in \bigcup_{n \in \mathbb{N}} \phi_n(\mathcal{B}_n) \subset \mathcal{B}$ and note that there exist $n \in \mathbb{N}$, $a \in \mathcal{B}_{n+1}$ and $b \in \mathcal{B}_n$ such that $\tilde{a} = \phi_{n+1}(a)$ and $\tilde{b} = \phi_n(b)$. Routine computation shows that $s_{*,n}(s_{n+1}(a)b) = a \cdot s_{*,n}(b)$ and thus using formulae (11) we have

$$\begin{aligned} \delta_*(\delta(\tilde{a})\tilde{b}) &= \delta_*(\phi_n(s_{n+1}(a))\tilde{b}) = \phi_{n+1}(s_{*,n}(s_{n+1}(a)b)) \\ &= \phi_{n+1}(a \cdot s_{*,n}(b)) = \phi_{n+1}(a) \cdot \phi_{n+1}(s_{*,n}(b)) = \tilde{a}\delta_*(\tilde{b}) \end{aligned}$$

what proves (2). Similarly we prove (3). It suffices to check that $s_{n+1}(s_{*,n}(a)) = s_{n+1}(1)as_{n+1}(1)$ and then we have

$$\begin{aligned} \delta(\delta_*(\tilde{a})) &= \delta(\phi_{n+1}(s_{*,n}(a))) = \phi_n(s_{n+1}(s_{*,n}(a))) = \phi_n(s_{n+1}(1)as_{n+1}(1)) \\ &= \phi_n(s_{n+1}(1))\phi_n(a)\phi_n(s_{n+1}(1)) = \delta(1)\tilde{a}\delta(1). \end{aligned}$$

■

The above presented construction of the system $(\mathcal{B}, \delta)$ is compatible with operator (geometric) approach presented in the introduction to this section.

Theorem 3.4. *There is a one-to-one correspondence between the strict covariant representations (π, U, H) and $(\tilde{\pi}, U, H)$ of $(\mathcal{A}, \delta)$ and $(\mathcal{B}, \delta)$ respectively, which is established by the relation*

$$\tilde{\pi}(\phi_n(a_0 \oplus a_1 \dots \oplus a_n)) = \pi(a_0) + U^* \pi(a_1)U \dots + U^{*n} \pi(a_n)U^n \quad (12)$$

Proof. The fact that for a strict covariant representation $(\tilde{\pi}, U, H)$ of $(\mathcal{B}, \delta)$ the triple (π, U, H) where $\pi = \tilde{\pi}|_{\mathcal{A}}$ is a covariant representation of $(\mathcal{A}, \delta)$ is trivial. However the strictness of (π, U, H) requires a comment. In particular, it follows from the equality of kernels of $\delta : \mathcal{A} \rightarrow \mathcal{A}$ and $\delta : \mathcal{B} \rightarrow \mathcal{B}$.

To prove the equality between $\ker \delta|_{\mathcal{B}}$ and $\ker \delta|_{\mathcal{A}} = q\mathcal{A}$ let $n > 0$ and notice that

$$\phi_0(q) = \phi_n(q \oplus 0 \oplus 0 \dots \oplus 0).$$

Thus $q\mathcal{B} = q\mathcal{A}$ and for $a = \phi_n(a_0 \oplus a_1 \oplus \dots \oplus a_n) \in \phi_n(\mathcal{B}_n)$ we have

$$\delta(a) = \phi_n(a_1 \oplus \dots \oplus a_n) = 0 \iff a_1 = \dots = a_n = 0 \iff a = qa.$$

Hence $\ker \delta|_{\mathcal{B}} = \ker \delta|_{\mathcal{A}}$.

Fix now a strict covariant representation (π, U, H) of $(\mathcal{A}, \delta)$. We show that formula (12) defines a faithful representation $\tilde{\pi}$ of $\mathcal{B}$.

The first step is to check that $\tilde{\pi}$ is well defined. Let $a \in \mathcal{B}_n$ and $b \in \mathcal{B}_m$ be such

that $\phi_n(a) = \phi_m(b)$ and let us assume that $n \leq m$ and $a = a_0 \oplus a_1 \oplus \dots \oplus a_n$. Using injectivity of the mappings occurring in the following equality

$$\phi_n = \phi_m \circ \delta_{m-1} \circ \dots \circ \delta_{n+1} \circ \delta_n$$

we obtain that b is of the form

$$b = a_0 \oplus \dots \oplus a_{n-1} \oplus qa_n \oplus q\delta(a_n) \oplus \dots \oplus q\delta^{m-n-1}(a_n) \oplus \delta^{m-n}(a_n).$$

Whereas covariance and strictness of (π, U, H) (Definition 1.6) together with the relation $a_n \in \mathcal{A}_n = \delta^n(1)\mathcal{A}\delta^n(1)$ implies that

$$\begin{aligned} U^{*n+k}\pi(q\delta^k(a_n))U^{n+k} &= U^{*n+k}(1 - U^*U)(U^k\pi(a_n)U^{*k})U^{n+k} \\ &= U^{*n+k}(U^k\pi(a_n)U^{*k})U^{n+k} - U^{*n+k}(U^*UU^k\pi(a_n)U^{*k})U^{n+k} \\ &= U^{*n}(U^{*k}U^k)\pi(a_n)(U^{*k}U^k)U^n - U^{*n}(U^{*(k+1)}U^{k+1}\pi(a_n)U^{*k}U^k)U^n \\ &= U^{*n}(U^{*k}U^k)\pi(a_n)U^n - U^{*n}(U^{*(k+1)}U^{k+1}\pi(a_n))U^n \\ &= U^{*n}((U^{*k}U^k - U^{*(k+1)}U^{k+1})\pi(a_n))U^n \\ &= (U^{*(n+k)}U^k - U^{*(n+k+1)}U^{k+1})\pi(\delta^n(1)a_n)U^n \\ &= (U^{*(n+k)}U^{n+k} - U^{*(n+k+1)}U^{n+k+1})U^{*n}\pi(a_n)U^n \end{aligned}$$

and

$$\begin{aligned} U^{*m}\pi(\delta^{m-n}(a_n))U^m &= U^{*m}U^{m-n}\pi(a_n)U^{*m-n}U^m \\ &= U^{*m}U^{m-n}(U^nU^{*n}\pi(a_n)U^nU^{*n})U^{*m-n}U^m = U^{*m}U^m(U^{*n}\pi(a_n)U^n)U^{*m}U^m \\ &= U^{*m}U^m(U^{*n}\pi(a_n)U^n) \end{aligned}$$

Thus we have

$$\sum_{k=0}^{m-n-1} U^{*n+k}\pi(Q\delta^k(a_n))U^{n+k} + U^{*m}\delta^{m-n}\pi((a_n))U^m = U^{*n}\pi(a_n)U^n.$$

Hence $\tilde{\pi}(\phi_n(a)) = \tilde{\pi}(\phi_m(b))$ and $\tilde{\pi}$ is well defined.

In view of Proposition 3.1 for every $n \in \mathbb{N}$ $\tilde{\pi} : \phi_n(\mathcal{B}_n) \rightarrow C^*(\bigcup_{k=0}^n U^{*n}\pi(\mathcal{A})U^n)$ is an isomorphism. Consequently $\tilde{\pi} : \mathcal{B} \rightarrow C^*(\bigcup_{n \in \mathbb{N}} U^{*n}\pi(\mathcal{A})U^n)$ is an isomorphism. Thus, the only thing left for us to show is that the $(\tilde{\pi}, U, H)$ satisfies conditions $(CR1'')$ and $(CR2)$. Condition $(CR1'')$ follows from the equality $\tilde{\pi}(q) = \pi(q) = U^*U$ and the coincidence of kernels $\ker \delta|_{\mathcal{B}} = \ker \delta|_{\mathcal{A}}$ (see the first part of the proof). To show $(CR2)$ take any element of the form $a = \phi_n(a_0 \oplus a_1 \oplus \dots \oplus a_n) \in \phi_n(\mathcal{B}_n)$, $n > 0$. Then

$$\begin{aligned} U\tilde{\pi}(a)U^* &= U\pi(a_0)U^* + UU^*\pi(a_1)UU^* \dots + UU^{*n}\pi(a_n)U^nU^* \\ &= \pi(\delta(a_0)) + \pi(\delta(1)a_1\delta(1)) \dots + UU^*(U^{*n-1}U^{n-1})U^{*n-1}\pi(a_n)U^{n-1}(U^{*n-1}U^{n-1})UU^* \\ &= \pi(a_1) \dots + U^{*n-1}U^nU^{*n}\pi(a_n)U^nU^{*n}U^{n-1} = \pi(a_1) \dots + U^{*n-1}\pi(\delta^n(1)a_n\delta^n(1))U^{n-1} \\ &= \pi(a_1) \dots + U^{*n-1}\pi(a_n)U^{n-1} = \tilde{\pi}(\phi_{n-1}(a_1 \oplus a_2 \oplus \dots \oplus a_n)) = \tilde{\pi}(\delta(a)). \end{aligned}$$

■

We put together the systems constructed in this and the previous subsection in a definition.

Definition 3.5. Let $(\mathcal{A}, \delta)$ be an arbitrary C^* -dynamical system. Let $(\mathcal{A}^+, \delta)$ be extension of $(\mathcal{A}, \delta)$ defined in the subsection 1.10 and let $(\mathcal{B}, \delta)$ be the extension of $(\mathcal{A}^+, \delta)$ described in the present subsection:

$$\mathcal{A} \subset \mathcal{A}^+ \subset \mathcal{B}.$$

We shall call the system $(\mathcal{B}, \delta)$ a *natural extension* of $(\mathcal{A}, \delta)$ to the C^* -dynamical system possessing complete transfer operator δ_* .

In view of Theorems 3.2 and 3.4 we have the relations

$$\mathcal{A}^+ = C^*(\mathcal{A}, \delta_*(1)), \quad \mathcal{B} = C^*\left(\bigcup_{n=0}^{\infty} \delta_*^n(\mathcal{A})\right),$$

where $\delta_* : \mathcal{B} \rightarrow \mathcal{B}$ is the transfer operator for $(\mathcal{B}, \delta)$, and we are ready to state the main result of the present paper.

Theorem 3.6. *Let $(\mathcal{A}, \delta)$ be an arbitrary C^* -dynamical system and let $(\mathcal{B}, \delta)$ be its natural extension with the transfer operator δ_* . There is a one-to-one correspondence between the strict covariant representations (π, U, H) and $(\tilde{\pi}, U, H)$ of $(\mathcal{A}, \delta)$ and $(\mathcal{B}, \delta)$ respectively, which is established by the relation*

$$\tilde{\pi}\left(\sum_{k=0}^n \delta_*^k(a_k)\right) = \sum_{k=0}^n U^{*k} \pi(a_k) U^k, \quad a_k \in \mathcal{A}.$$

In particular for every strict covariant representation (π, U, H) of $(\mathcal{A}, \delta)$ we have

$$\mathcal{B} \cong C^*\left(\bigcup_{n \in \mathbb{N}} U^{*n} \pi(\mathcal{A}) U^n\right).$$

We conclude this section with the nontrivial fact which generalizes the essential part of the main result of [BL05]. It is a corollary to Theorems 3.6, 2.5 and 2.6.

Theorem 3.7. *For an arbitrary C^* -dynamical system $(\mathcal{A}, \delta)$ there exists a strict covariant representation of $(\mathcal{A}, \delta)$.*

4 Commutative case and its relation to inverse limits

Throughout this section $\mathcal{A}$ is a unital commutative C^* -algebra. Let us recall that the space X of all non-zero linear and multiplicative functionals on $\mathcal{A}$ equipped with the weak* topology is a Hausdorff space and by Gelfand-Naimark Theorem $\mathcal{A}$ is isomorphic to (and thus may be identified with) the algebra $C(X)$ of continuous complex valued functions on X . It is also known, cf. [Bou67], that then an endomorphism $\delta : \mathcal{A} \rightarrow \mathcal{A}$ assumes the form

$$\delta(a)(x) = \begin{cases} a(\alpha(x)) & , \quad x \in \Delta_1 \\ 0 & , \quad x \notin \Delta_1 \end{cases}, \quad a \in C(X), \quad (13)$$

where Δ_1 is a clopen subset of X and $\alpha : \Delta_1 \rightarrow X$ is a continuous map. We will denote by $\Delta_{-1} = \alpha(\Delta_1)$ the image of α and refer to (X, α) as to *partial dynamical system* generated by $(\mathcal{A}, \delta)$.

We have a number of dual correspondences between the properties of C^* -dynamical systems and partial systems they generate, cf. [KL03], [Kwa07]. Let us list a few of them:

$$\begin{array}{ll}
 \delta \text{ is a monomorphism} & \iff \alpha \text{ is a surjection,} \\
 \delta \text{ is unital} & \iff \text{the domain of } \alpha \text{ is } X, \\
 \delta \text{ is an epimorphism} & \iff \alpha \text{ is an injection with domain } X, \\
 \text{the kernel of } \delta \text{ is unital} & \iff \text{the image of } \alpha \text{ is open,} \\
 \left(\begin{array}{l} \text{the system } (\mathcal{A}, \delta) \text{ possess} \\ \text{a complete transfer operator} \end{array} \right) & \iff \left(\begin{array}{l} \alpha \text{ is an injection and} \\ \text{the image of } \alpha \text{ is open} \end{array} \right).
 \end{array}$$

This specific spectral duality allows us to explore the constructions from previous section in topological terms.

4.1 Compactification of a complement of a range of a mapping

Adjoining a unit to a non-unital commutative algebra corresponds to compactification of a non-compact topological space. The kernel I of the endomorphism $\delta : \mathcal{A} \rightarrow \mathcal{A}$ is isomorphic to the algebra $C_0(X \setminus \Delta_{-1})$. Hence adjoining a unit to I corresponds to a compactification of $X \setminus \Delta_{-1}$. The construction of the system $(\mathcal{A}^+, \delta)$ from subsection 3.1 relies simply on taking the closure $\overline{X \setminus \Delta_{-1}}$ of $X \setminus \Delta_{-1}$ in X .

Indeed, the system $(\mathcal{A}^+, \delta)$ is commutative and thereby it generates a partial dynamical (X_+, α_+) which may be informally described in terms of (X, α) in the following way, see Figure 4.1: the space X_+ arise by "cutting" Δ_{-1} out of X and then closing the set $X \setminus \Delta_{-1}$; the image of α_+ is the set Δ_{-1} that has been "cut out" and the domain of α_+ , with respect to the domain of α , is enlarged by the set $\Delta_1 \cap \overline{X \setminus \Delta_{-1}}$. The whole space X_+ differs from X by a boundary $\text{Bd}(\Delta_{-1}) = \Delta_{-1} \cap \overline{X \setminus \Delta_{-1}}$ of the range Δ_{-1} of α .




Figure 1: Compactification of a complement of a range of a mapping.

Formally, X_+ is the direct sum of two topological spaces X_1 and X_2 homeomorphic

to $\overline{X \setminus \Delta_{-1}}$ and Δ_{-1} respectively:

$$\mathcal{A}/I = A/C_0(X \setminus \Delta_{-1}) = C(X_1) \cong C(\Delta_{-1}), \quad \mathcal{A}/I^\perp = C(X_2) \cong C(\overline{X \setminus \Delta_{-1}}).$$

If we let $\phi_1 : X_1 \rightarrow \Delta_{-1}$ and $\phi_2 : X_2 \rightarrow \overline{X \setminus \Delta_{-1}}$ be the corresponding homeomorphisms then the mapping α_+ is defined on the set

$$\phi_1^{-1}(\Delta_1) \cup \phi_2^{-1}(\Delta_1)$$

by the formula

$$\alpha_+(x) = \begin{cases} \phi_2^{-1}(\alpha(\phi_1(x))), & x \in \phi_1^{-1}(\Delta_1) \\ \phi_2^{-1}(\alpha(\phi_2(x))), & x \in \phi_2^{-1}(\Delta_1) \end{cases}.$$

4.2 Inverse limits and spectral description

It is a straightforward fact, cf. [LO04, Proposition 4.1], that the natural extension $(\mathcal{B}, \delta)$ (see Definition 3.5) of the commutative system $(\mathcal{A}, \delta)$ is also commutative. A description of the partial dynamical system generated by $(\mathcal{B}, \delta)$ was the purpose of [KL03]. The method elaborated there differs and has certain advantages over the general setting included in subsection 3.2 (the difference is discussed in the coming subsection). Here, we slightly improve the main result of [KL03] and clarify its relation to inverse limits.

To start with we recall the definition of inverse limits in the category of topological spaces.

Definition 4.1. *An inverse limit of an inverse sequence*

$$X_0 \xleftarrow{\alpha_0} X_1 \xleftarrow{\alpha_1} X_2 \xleftarrow{\alpha_2} \dots$$

where X_n are topological spaces and $\alpha_n : X_{n+1} \rightarrow X_n$ are continuous maps is the topological space of the form

$$\varprojlim \{X_n, \alpha_n\} = \{(x_0, x_1, \dots) \in \prod_{n \in \mathbb{N}} X_n : \alpha_n(x_{n+1}) = x_n, n \in \mathbb{N}\}$$

considered with the product topology inherited from $\prod_{n \in \mathbb{N}} X_n$. If additionally all the spaces and mappings coincide, that is if for every $n \in \mathbb{N}$ we have $X_n = X$ and $\alpha_n = \alpha$, then we write $\varprojlim \{X_n, \alpha_n\} = \varprojlim \{X, \alpha\}$ and the formula

$$\tilde{\alpha}(x_0, x_1, \dots) = (\alpha(x_0), x_0, x_1, \dots), \quad (x_0, x_1, \dots) \in \varprojlim \{X, \alpha\},$$

defines a homeomorphism $\tilde{\alpha} : \varprojlim \{X, \alpha\} \rightarrow \varprojlim \{X, \alpha\}$. We say that $\tilde{\alpha}$ is *induced* by $\alpha : X \rightarrow X$.

Now, let (X, α) be the partial dynamical system generated by $(\mathcal{A}, \delta)$ and let $(\tilde{X}, \tilde{\alpha})$ be the partial dynamical system generated by the natural extension $(\mathcal{B}, \delta)$ of $(\mathcal{A}, \delta)$. We extend our notation and for any $n \in \mathbb{N}$ we denote by $\Delta_n = \alpha^{-n}(X)$ the

domain of α^n and by $\Delta_{-n} = \alpha^n(\Delta_n)$ its image. A similar "tilde-notation" will be used for $\tilde{\alpha}$. Thus we have

$$\alpha^n : \Delta_n \rightarrow \Delta_{-n} \quad \text{and} \quad \tilde{\alpha}^n : \tilde{\Delta}_n \rightarrow \tilde{\Delta}_{-n}.$$

Moreover, $\tilde{\alpha} : \tilde{\Delta}_1 \rightarrow \tilde{\Delta}_{-1}$ is a homeomorphism and the complete transfer operator $\delta_* : \mathcal{B} \rightarrow \mathcal{B}$ for $(\mathcal{B}, \delta)$ is an endomorphism of the form

$$\delta_*(a)(\tilde{x}) = \begin{cases} a(\tilde{\alpha}^{-1}(\tilde{x})) & , \quad \tilde{x} \in \tilde{\Delta}_{-1} \\ 0 & , \quad \tilde{x} \notin \tilde{\Delta}_{-1} \end{cases}, \quad a \in \mathcal{B} = C(\tilde{X}). \quad (14)$$

In order to establish the relationship between the systems (X, α) and $(\tilde{X}, \tilde{\alpha})$ we fix for a while an arbitrary point $\tilde{x} \in \tilde{X}$. Then considering $\tilde{x}$ as a linear and multiplicative functional on $\mathcal{B}$ we obtain a sequence of functionals $x_n : \mathcal{A} \rightarrow \mathbb{C}$, $n = 0, 1, \dots$, defined by the condition

$$x_n(a) = \tilde{x}(\delta_*^n(a)), \quad a \in \mathcal{A}.$$

Since the set $\bigcup_{n=0}^{\infty} \delta_*^n(\mathcal{A})$ is linearly dense in $\mathcal{B}$, the sequence $(x_0, x_1, \dots)$ determines $\tilde{x}$ uniquely. Multiplicativity of δ_* on $\mathcal{B}$ implies that all functionals $\tilde{x}_n$ are linear and multiplicative, hence

$$\text{either } x_n \in X \quad \text{or} \quad x_n \equiv 0.$$

Summing it all up the mapping

$$\tilde{X} \ni \tilde{x} \longrightarrow (x_0, x_1, x_2, \dots) \in \prod_{n=0}^{\infty} X \cup \{0\} \quad (15)$$

is injective and under the assumption that the set Δ_{-1} is open (equivalently kernel of $\delta : \mathcal{A} \rightarrow \mathcal{A}$ is unital) the authors of [KL03] proved the following

Theorem 4.2. *If Δ_{-1} is open, then mapping (15) defines a homeomorphism under which we have the following identifications*

$$\tilde{X} = \bigcup_{N=0}^{\infty} X_N \cup X_{\infty} \quad (16)$$

where

$$X_N = \{(x_0, x_1, \dots, x_N, 0, \dots) : x_n \in \Delta_n, \alpha(x_n) = x_{n-1}, n = 1, \dots, N, x_N \notin \Delta_{-1}\},$$

$$X_{\infty} = \{(x_0, x_1, \dots) : x_n \in \Delta_n, \alpha(x_n) = x_{n-1}, n \geq 1\},$$

$\bigcup_{N=0}^{\infty} X_N \cup X_{\infty}$ is equipped with the product topology inherited from $\prod_{n=0}^{\infty} X \cup \{0\}$ (set $\{0\}$ is clopen in $X \cup \{0\}$). The partial homeomorphism $\tilde{\alpha}$ is of the form

$$\tilde{\alpha}(x_0, \dots) = (\alpha(x_0), x_0, \dots), \quad x \in \tilde{\Delta}_1,$$

where $\tilde{\Delta}_1 = \{(x_0, \dots) \in \tilde{X} : x_0 \in \Delta_1\}$ is the domain of $\tilde{\alpha}$.

We notice that the subset $X_\infty \subset \tilde{X}$ is the inverse limit of the following inverse sequence

$$X \xleftarrow{\alpha} \Delta_1 \xleftarrow{\alpha} \Delta_2 \xleftarrow{\alpha} \dots$$

Elements of X_N may be considered as finite anty-orbits of α which cannot be prolonged (because $x_N \notin \Delta_{-1}$).

In view of Definition 3.5 and subsection 4.1 one can generalize Theorem 4.2 to get a result without the assumption of openness of Δ_{-1} . Roughly speaking, it suffices to replace (in the description of sets X_N) the condition $x_N \notin \Delta_{-1}$ with a weaker one $x_N \in \overline{X \setminus \Delta_{-1}}$. More formally one needs to take the closure of $\tilde{X}$ in $\prod_{n=0}^{\infty} X \cup \{0\}$. We shall express this remark in a definition and a theorem.

Definition 4.3. Let (X, α) be an arbitrary partial dynamical system, that is $\alpha : \Delta_1 \rightarrow X$ is a continuous mapping and Δ_{-1} is a clopen subset of a compact space X . By a *natural extension* of (X, α) we shall mean the partial dynamical system $(\tilde{X}, \tilde{\alpha})$ where

$$\tilde{X} = \bigcup_{N=0}^{\infty} X_N \cup X_\infty,$$

$$X_N = \{(x_0, x_1, \dots, x_N, 0, \dots) : x_n \in \Delta_n, \alpha(x_n) = x_{n-1}, n = 1, \dots, N, x_N \in \overline{X \setminus \Delta_{-1}}\},$$

$$X_\infty = \{(x_0, x_1, \dots) : x_n \in \Delta_n, \alpha(x_n) = x_{n-1}, n \geq 1\},$$

is equipped with the product topology $\prod_{n=0}^{\infty} X \cup \{0\}$, set $\{0\}$ is clopen and 0 is just an abstract point. Mapping $\tilde{\alpha} : \tilde{\Delta}_1 \rightarrow \tilde{\Delta}_{-1}$ is a homeomorphism given by the formula

$$\tilde{\alpha}(x_0, \dots) = (\alpha(x_0), x_0, \dots), \quad x \in \tilde{\Delta}_1, \quad (17)$$

where $\tilde{\Delta}_1 = \{(x_0, \dots) \in \tilde{X} : x_0 \in \Delta_1\}$ and $\tilde{\Delta}_{-1} = \{(x_0, x_1, \dots) \in \tilde{X} : x_1 \in \Delta_1\}$.

The notion of natural extension for topological and for C^* -dynamical systems agrees in the following sense.

Theorem 4.4. *The natural extension $(\tilde{X}, \tilde{\alpha})$ of a partial dynamical system (X, α) is the partial dynamical system generated by the natural extension $(\mathcal{B}, \delta)$ of the C^* -dynamical system $(\mathcal{A}, \delta)$ that generates (X, α) .*

Proof. It is a corollary to Definition 3.5, subsection 4.1 and Theorem 4.2. ■

Let us now consider the classical case, that is when mapping $\alpha : X \rightarrow X$ is described on the whole space X . Then the natural extension $(\tilde{X}, \tilde{\alpha})$ has the following structure:

- the system $(X_\infty, \tilde{\alpha})$ coincide with the inverse limit system $(\varprojlim \{X, \alpha\}, \tilde{\alpha})$,
- for every $N \in \mathbb{N}$ the set X_N is homeomorphic to $\overline{X \setminus \Delta_{-1}}$ and $\tilde{\alpha}$ maps homeomorphically X_N onto X_{N+1} :

$$X_0 \xrightarrow{\tilde{\alpha}} X_1 \xrightarrow{\tilde{\alpha}} X_2 \xrightarrow{\tilde{\alpha}} \dots (X_\infty, \tilde{\alpha}) = (\varprojlim \{X, \alpha\}, \tilde{\alpha}).$$

In particular the system $(\tilde{X}, \tilde{\alpha})$ is not a direct generalization of the inverse limit system $(\varprojlim \{X, \alpha\}, \tilde{\alpha})$, it is rather an "improvement" of it, as it contains more information concerning the initial system (X, α) .

Example 4.5. Let $\alpha : X \rightarrow X$ be a constant mapping with the only value being a non-isolated point $p \in X$.




Figure 2: Systems associated with constant map.

Then the system (X_+, α_+) from paragraph 4.1 is obtained from (X, α) by "cutting out" the point $\{p\}$ and "filling the hole that arose during this operation". Space $\tilde{X}$ is the countable number of copies of X compactified with one point $X_\infty = \{p\}$, whereas the system $(\varprojlim \{X, \alpha\}, \tilde{\alpha})$ degenerates to a single point, see Figure 2.

4.3 Comparison of the spectral and the algebraic approach

Description of the system $(\tilde{X}, \tilde{\alpha})$ presented in Theorem 4.2 and description of $(\mathcal{B}, \delta)$ presented in Theorem 3.3 (hence also of $(\tilde{X}, \tilde{\alpha})$ in the commutative case) are obtained by different methods. As the relationship between those construction is not obvious and might be interesting we include a short discussion on the subject.

We assume here that Δ_{-1} is open (equivalently $\overline{X \setminus \Delta_{-1}}$ is closed). This assumption is not necessary but it simplifies notation. Replacing every set of the form $K \setminus \Delta_{-1}$ with a bigger one of the form $K \cap \overline{X \setminus \Delta_{-1}}$ everything what follows remains true in general.

We recall that algebra $\mathcal{B} \cong C(\tilde{X})$ is the direct limit of the direct sequence (8) with bonding homomorphisms δ_n that are unital monomorphisms, see subsection 3.2. Thus the space $\tilde{X} = \varprojlim \{\tilde{X}_n, \alpha_n\}$ is the inverse limit of the inverse limit

$$\tilde{X}_0 \xleftarrow{\alpha_0} \tilde{X}_1 \xleftarrow{\alpha_1} \tilde{X}_2 \xleftarrow{\alpha_2} \dots \quad (18)$$

where $\mathcal{B}_n = C(\tilde{X}_n)$ and $\alpha_n : \tilde{X}_{n+1} \rightarrow \tilde{X}_n$ are surjections such that, $\delta_n(a)(x) = a(\alpha_n(x))$, $a \in \mathcal{B}_n$. From definition (7) of $\mathcal{B}_n$ we have $\tilde{X}_n$ is a direct sum of clopen subsets of X . Namely, using the symbol " $\sqcup$ " to denote the disjoint union we get

$$\tilde{X}_n = (X \setminus \Delta_{-1}) \sqcup (\Delta_1 \setminus \Delta_{-1}) \sqcup \dots \sqcup (\Delta_{n-1} \setminus \Delta_{-1}) \sqcup \Delta_n.$$

The inverse sequence (18) maybe rewritten in the form of "triangular" diagram illustrating also the action of the mappings $\alpha_n : \tilde{X}_{n+1} \rightarrow \tilde{X}_n$:

$$\begin{array}{rcl}
 \tilde{X}_0 & = & X \\
 \uparrow \alpha_1 & & \uparrow id \quad \swarrow \alpha \\
 \tilde{X}_1 & = & X \setminus \Delta_{-1} \sqcup \Delta_1 \\
 \uparrow \alpha_2 & & \uparrow id \quad \uparrow id \quad \swarrow \alpha \\
 \tilde{X}_2 & = & X \setminus \Delta_{-1} \sqcup \Delta_1 \setminus \Delta_1 \sqcup \Delta_2 \\
 \uparrow \alpha_3 & & \uparrow id \quad \uparrow id \quad \uparrow id \quad \swarrow \alpha \\
 \tilde{X}_3 & = & X \setminus \Delta_{-1} \sqcup \Delta_1 \setminus \Delta_1 \sqcup \Delta_2 \setminus \Delta_{-1} \sqcup \Delta_3 \\
 \vdots & & \vdots \quad \vdots \quad \vdots \quad \vdots \quad \ddots
 \end{array}$$

Thus, if $\tilde{x} = (x_0, x_1, x_2, \dots)$ is an element of the inverse limit $\tilde{X} = \varprojlim \{\tilde{X}_n, \alpha_n\}$, then we have two alternatives: either every coordinate of $\tilde{x}$ "lies on the diagonal" of the above diagram, that is $\tilde{x}$ belongs to the set of the form

$$X_\infty = \{(x_0, x_1, \dots) \in X^\mathbb{N} : x_n \in \Delta_n, \alpha(x_n) = x_{n-1}, n > 0\},$$

or only N first coordinates of $\tilde{x}$ "lies on the diagonal" and next ones "slide off the diagonal and stays out of it forever", that is $\tilde{x}$ belongs to the set of the form

$$X_N = \{(x_0, x_1, \dots, x_N, 0, 0, \dots) : x_n \in \Delta_n, \alpha(x_n) = x_{n-1}, n = 1, \dots, N, x_N \notin \Delta_{-1}\},$$

where we replaced the tail of x_N 's with tail of zeros.

It is not hard to see that the above considerations lead us to identification of the inverse limit $\tilde{X} = \varprojlim \{\tilde{X}_n, \alpha_n\}$ with the space $\bigcup_{n \in \mathbb{N}} X_n \cup X_\infty$ from Theorem 4.2. It is a bit harder to realize the relationship between formula (17) and inverse sequence (9) of homomorphisms $s_{n+1} : \mathcal{B}_{n+1} \rightarrow \mathcal{B}_n$. In fact, a homomorphism s_{n+1} induces a continuous mapping defined on the clopen subset of $\tilde{X}_n$ with values in $\tilde{X}_{n+1}$, given by the following diagram:

$$\begin{array}{rcl}
 \tilde{X}_n & \supset & \Delta_1 \setminus \Delta_{-1} \sqcup \Delta_2 \setminus \Delta_{-1} \sqcup \dots \sqcup \Delta_{n+1} \\
 & & \searrow id \quad \searrow id \quad \searrow id \\
 \tilde{X}_{n+1} & = & X \setminus \Delta_{-1} \sqcup \Delta_1 \setminus \Delta_{-1} \sqcup \Delta_2 \setminus \Delta_{-1} \sqcup \dots \sqcup \Delta_{n+1}
 \end{array}$$

Then one needs to check that the direct sequence of the above diagrams defines a mapping on a clopen subset of the inverse limit of the inverse sequence (18).

This "tangle" of direct and inverse sequences which arise in the general approach proves the usefulness of the spectral description from Theorems 4.2, 4.4 which simply presents the system $(\tilde{X}, \tilde{\alpha})$ as a shift on a certain space of sequences.

References

- [AP98] J. E. Anderson, I. F. Putnam, "Topological invariants for substitution tilings and their associated C^* -algebras", *Ergod. Th. & Dynam. Sys.*, **18**, (1998), pp. 509–537.
- [ABL05] A.B. Antonevich, V.I. Bakhtin, A.V. Lebedev, "Crossed product of C^* -algebra by an endomorphism, coefficient algebras and transfer operators", arXiv:math.OA/0502415 v1 19 Feb 2005.
- [BL05] V.I. Bakhtin, A.V. Lebedev, "When a C^* -algebra is a coefficient algebra for a given endomorphism", arXiv:math.OA/0502414 v1 19 Feb 2005.
- [Bou67] N. Bourbaki *Théories spectrales*, Herman 1967.
- [CK80] J. Cuntz and W. Krieger, "A Class of C^* -algebras and Topological Markov Chains", *Inventiones Math.* **56**, (1980), p. 251–268
- [Exel94] R. Exel, *Circle actions on C^* -algebras, partial automorphisms and generalized Pimsner-Voiculescu exact sequence*, *J. Funct. Analysis* **122** (1994), p. 361–401.
- [Exel03] R. Exel, "A new look at the crossed-product of a C^* -algebra by an endomorphism", *Ergodic Theory Dynam. Systems*, Vol 23, (2003), pp. 1733–1750,
- [KR86] Kadison R.V., Ringrose J.R., *Fundamentals of the theory of operator algebras. Vol.2. Advanced theory*, Academic Press, 1986.
- [Kwa05a] B.K. Kwaśniewski, "Covariance algebra of a partial dynamical system", *CEJM*, 2005, V.3, No 4, pp. 718–765, arXiv:math.OA/0407352
- [Kwa05b] B. K. Kwaśniewski, "Inverse limit systems associated with F_2^n zero schwarzian unimodal mappings", *Bull. Soc. Sci. Lett. Lodz*, Vol 55 (2005), pp. 83–109, arXiv:math.DS/0502584
- [Kwa07] B.K. Kwaśniewski: "On transfer operators for C^* -dynamical systems", preprint arXiv:math.OA/0703798 v1 27 Mar 2007
- [KL03] B.K. Kwaśniewski and A.V. Lebedev: "Maximal ideal space of a commutative coefficient algebra", preprint arXiv:math.OA/0311416
- [LO04] A.V. Lebedev, A. Odziejewicz "Extensions of C^* -algebras by partial isometries", *Matemat. Sbornik*, 2004. V. 195, No 7, pp. 37–70 (Russian).
- [Mur90] G. J. Murphy: *C^* -algebras and operator theory*, Academic Press, 1990.
- [Mur96] G. J. Murphy, "Crossed products of C^* -algebras by endomorphisms", *Integral Equations Oper. Theory* **24**, (1996), p. 298–319.
- [Nad92] S. Nadler, *Continuum Theory*, Dekker, New York, 1992.

- [Pas80] W.L. Paschke, "The crossed product of a C^* -algebra by an endomorphism", *Proceedings of the AMS*, **80**, No 1, (1980), p. 113–118.
- [Ped79] G. K. Pedersen, *C^* -algebras and their automorphism groups*, Academic Press, London, 1979.
- [Sak71] S. Sakai, *C^* -algebras and W^* -algebras*, Springer-Verlag, New-York, (1971).
- [Sta93] P. J. Stacey, "Crossed products of C^* -algebras by $*$ -endomorphisms", *J. Austral. Math. Soc. Ser. A* **54**, (1993), p. 204–212.
- [Wil67] R. F. Williams, "One-dimensional nonwandering sets", *Topology*, **6**, (1967), pp. 473–487.
- [Wil70] R. F. Williams, "Classification of 1-dimensional attractors", *Proc. Sym. Pure Math. AMS*, **14**, (1970), pp. 341–361.